

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/02090 A2

- (51) International Patent Classification⁷: **A61K 31/00**
- (21) International Application Number: **PCT/FI01/00638**
- (22) International Filing Date: **4 July 2001 (04.07.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/216,341 **5 July 2000 (05.07.2000)** **US**
- (71) Applicant (for all designated States except US): **BIOTIE THERAPIES CORP.** [FI/FI]; Tykistökatu 6, FIN-20520 Turku (FI).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **SMITH, David, John** [GB/FI]; Valliuksenkatu 6, FIN-21100 Naantali (FI). **JALKANEN, Markku** [FI/FI]; Rauvolantie, FIN-20760 Piispanristi (FI). **FÜLÖP, Ferenc** [HU/HU]; Petőfi S. sgt. 7, H-6722 Szeged (HU). **LÁZÁR, László** [HU/HU]; Alkony u. 4/A, H-6725 Szeged (HU). **SZAKONYI, Zsolt** [HU/HU]; Ortutay u. 6/A, H-6723 Szeged (HU). **BERNÁTH, Gábor** [HU/HU]; Merey u. 8, H-6722 Szeged (HU).
- (74) Agent: **OY JALO ANT-WUORINEN AB**; Iso Roobertinkatu 4-6 A, FIN-00120 Helsinki (FI).
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO** patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), **Eurasian** patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), **European** patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), **OAPI** patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **INHIBITORS OF COPPER-CONTAINING AMINE OXIDASES**

(57) Abstract: The present invention is directed to hydrazino compounds that functions as inhibitors of copper-containing amine oxidases commonly known as semicarbazidesensitive amine oxidases (SSAO), including the human SSAO known as Vascular Adhesion Protein-1 (VAP-1). These SSAO inhibitors have therapeutic utility as drugs to treat conditions and diseases including, but not limited to, a number of inflammatory conditions and diseases (in particular chronic inflammatory conditions such as chronic arthritis, inflammatory bowel diseases, and chronic skin dermatoses), diseases related to carbohydrate metabolism and to aberrations in adipocyte differentiation or function and smooth muscle cell function, and vascular diseases. The compounds have general formula: or a pharmaceutically acceptable solvate, hydrate, or salt thereof, wherein R¹ to R⁸ and X are as defined herein.

WO 02/02090 A2

Inhibitors Of Copper-Containing Amine Oxidases

5

Field of the Invention

10 The present invention is in the field of medicinal chemistry and is directed to hydrazino compounds and their use as inhibitors of copper-containing amine oxidases (E.C. 1.4.3.6) and enzymes of significant identity thereto. The compounds of the present invention have therapeutic utility as drugs to treat diseases including, but not limited to, inflammatory diseases. In particular, acute
15 and chronic inflammatory conditions or diseases such as chronic arthritis, inflammatory bowel diseases and skin dermatoses as well as diseases related to carbohydrate metabolism and to aberrations in adipocyte differentiation or function and smooth muscle cell function may be treated with the compounds.


20 *Background of the invention*

 VAP-1 is a human endothelial cell adhesion molecule that has several unique properties that distinguish it from the other inflammation-related adhesion molecules. It has a unique and restricted expression pattern and mediates
25 lymphocyte binding to vascular endothelium (Salmi, M., and Jalkanen, S., *Science* 257:1407-1409 (1992)). Inflammation induces the upregulation of VAP-1 to the surface of vascular endothelial cells mediating leukocyte entry to skin, gut and inflamed synovium (Salmi, M., and Jalkanen, S., *Science* 257:1407-1409 (1992); Salmi, M., *et al.*, *J. Exp. Med* 178:2255-2260 (1993); Arvillomi, A., *et al.*, *Eur. J.*
30 *Immunol.* 26:825-833 (1996); Salmi, M., *et al.*, *J. Clin. Invest.* 99:2165-2172 (1997); (Salmi, M., and Jalkanen, S., *J. Exp. Med.* 183:569-579 (1996); *J. Exp. Med.* 186:589-600 (1997)). One of the most interesting features of VAP-1 is a

- 2 -

1 catalytic extracellular domain which contains a monoamine oxidase activity
(Smith, D.J., *et al.*, *J. Exp. Med.* 188:17-27 (1998)).

2 The cloning and sequencing of the human VAP-1 cDNA revealed that it
encodes a transmembrane protein with homology to a class of enzymes called the
5 copper-containing amine oxidases (E.C. 1.4.3.6). Enzyme assays have shown that
VAP-1 possesses a monoamine oxidase (MAO) activity which is present in the
extracellular domain of the protein (Smith, D.J., *et al.*, *J. Exp. Med.* 188:17-27
(1998)). Thus, VAP-1 is an ecto-enzyme. Analysis of the VAP-1 MAO activity
10 showed that VAP-1 belongs to the class of membrane-bound MAO's termed
semicarbazide-sensitive amine oxidases (SSAO). These are distinguished from the
widely distributed mitochondrial MAO-A and B flavoproteins by amino acid
sequence, cofactor, substrate specificity and sensitivity to certain inhibitors.
However, certain substrates and inhibitors are common to both SSAO and MAO
activities. The mammalian SSAO's can metabolize various monoamines produced
15 endogenously or absorbed as dietary or xenobiotic substances. They act
principally on primary aliphatic or aromatic monoamines such as methylamine or
benzylamine (Lyles, G.A., *Int. J. Biochem. Cell Biol.* 28:259-274 (1996)). Thus,
VAP-1 located on the vascular endothelial cell surface can act on circulating
primary monoamines with the following reaction pathway.



The physiological substrates of VAP-1 SSAO in man have not been clearly
identified however methylamine is a good substrate for VAP-1 SSAO.
Methylamine is a product of various human biochemical pathways for the
25 degradation of creatinine, sarcosine and adrenaline, and is found in various
mammalian tissues and in blood. It can also be derived from the diet by gut
bacterial degradation of dietary precursors. The concentration of methylamine in
the blood can be increased in certain physiological and pathological situations such
as diabetes. Another potential physiological substrates is aminoacetone.

30 VAP-1 SSAO activity has been proposed to be directly involved in the
pathway of leukocyte adhesion to endothelial cells by a novel mechanism

- 3 -

involving direct interaction with an amine substrate presented on a VAP-1 ligand expressed on the surface of a leukocyte (Salmi *et al. Immunity*, (2001)). This publication describes the direct involvement of VAP-1 SSAO activity in the process of adhesion of leukocytes to endothelium. Thus inhibitors of VAP-1
5 SSAO activity could be expected to reduce leukocyte adhesion in areas of inflammation and thereby reduce leukocyte trafficking into the inflamed region and therefore the inflammatory process itself.


In human clinical tissue samples expression of VAP-1 is induced at sites of inflammation. This increased level of VAP-1 can lead to increased production of
10 H₂O₂ generated from the action of the VAP-1 SSAO extracellular domain on monoamines present in the blood. This generation of H₂O₂ in the localised environment of the endothelial cell could initiate other cellular events. H₂O₂ is a known signalling molecule that can upregulate other adhesion molecules and this increased adhesion molecule expression may lead to enhanced leukocyte
15 trafficking into areas in which VAP-1 is expressed. It also may be that other products of the VAP-1 SSAO reaction could have biological effects also contributing to the inflammatory process. Thus the products of the VAP-1 SSAO activity may be involved in an escalation of the inflammatory process which could be blocked by specific SSAO inhibitors.

20 VAP-1 SSAO may be involved in a number of other pathological conditions associated with an increased level of circulating amine substrates of VAP-1 SSAO. The oxidative deamination of these substrates would lead to an increase in the level of toxic aldehydes and oxygen radicals in the local environment of the endothelial cell which could damage the cells leading to
25 vascular damage. Increased levels of methylamine and aminoacetone have been reported in patients with Type I and Type II diabetes and it has been proposed that the vasculopathies such as retinopathy, neuropathy and nephropathy seen in late stage diabetes could be treated with specific inhibitors of SSAO activity.

Takahashi, H., *et al.*, *Yakugaku Zasshi* 101(12):1154-1156 (1981), report
30 the synthesis of a number of *N*-alkylaminoephedrine, including *N*-


- 4 -

(isopropylideneamino)-ephedrine (or *R,S*-(+)-(2-hydroxy-1-methyl-2-phenylethyl)methylhydrazone-2-propanone):



These hydrazone compounds were synthesized to evaluate their effect on the
5 bronchial musculature and were found not to exhibit any significant activity.

Grifantini, M., *et al.*, *Farmaco, Ed. Sci.* 23(3):197-203 (1968), report the
synthesis of several alkyl- and acyl-derivatives of *N*-amino-1-ephedrine and *N*-
amino-*d*-pseudoephedrine having antidepressant and monoamine oxidase
inhibitory properties. Among the compounds disclosed is the hydrazone erythro-
10 (β-hydroxy-α-methylphenethyl)methylhydrazone cyclohexanone, which has the
following structure:



The development of specific VAP-1 SSAO inhibitors that modulate VAP-1
activity would be useful for the treatment of acute and chronic inflammatory
15 conditions or diseases such as chronic arthritis, inflammatory bowel diseases, and
skin dermatoses, as well as diseases related to carbohydrate metabolism (including
diabetes and complications resulting from diabetes). In addition, aberrations in
adipocyte differentiation or function and smooth muscle cell function (in
particular, atherosclerosis), and various vascular diseases may be suitable for
20 treatment with VAP-1 SSAO inhibitors.

Summary of the Invention

5 The present invention is broadly directed to the use of hydrazino
compounds of Formula I or II as inhibitors of the class of copper-containing amine
oxidases known as semicarbazide-sensitive amine oxidases (SSAO), including the
human SSAO known as Vascular Adhesion Protein-1 (VAP-1). As VAP-1 SSAO
inhibitors, compounds of the present invention can function to prevent leukocyte
10 VAP-1 SSAO. Compounds of the present invention are therefore useful for
treating a number of inflammatory conditions and diseases of connective tissue,
skin, and the gastrointestinal, central nervous system, and pulmonary systems,
including such conditions as chronic arthritis, inflammatory bowel diseases, and
chronic dermatoses. The compounds are also useful for treating diseases related
15 to carbohydrate metabolism (such as diabetes), to aberrations in adipocyte
differentiation or function or smooth muscle cell function (such as atherosclerosis
and obesity), and to various vascular diseases (such as atheromatous and
nonatheromatous atherosclerosis, ischemic heart disease, and peripheral arterial
occlusion).

20 A further aspect of the present invention is to provide a pharmaceutical
composition useful for treating disorders responsive to a decrease in SSAO
activity, containing an effective amount of a compound of Formula I or II in a
mixture with one or more pharmaceutically acceptable carriers or diluents.


- 6 -

Detailed Description of the Invention

One aspect of the invention is to use a specific group of hydrazino compounds having the general formula I or II as defined below, for the manufacture of a pharmaceutical preparation for inhibiting a copper-containing amine oxidase.

Another aspect of the invention is to use a specific group of hydrazino compounds having the general formula I or II as defined below, for the manufacture of a pharmaceutical preparation for the treatment of an inflammatory disease or condition, a disease related to carbohydrate metabolism, a disease related to aberrations in adipocyte differentiation or function or smooth muscle cell function, or a vascular disease.

A further aspect of the present invention is directed to a method of inhibiting a copper-containing amine oxidase, the method comprising contacting said amine oxidase with an inhibitory effective amount of a hydrazino compound of Formula I or II either in racemates or optically pure forms



as a racemate or an isomer, or a pharmaceutically acceptable solvate, hydrate, or salt thereof;

wherein:

R¹ is hydrogen, (C₁-C₄)alkyl, aralkyl, (C₂-C₅)alkanoyl, aroyl or heteroaroyl,

R² is hydrogen, or optionally substituted (C₁-C₄)alkyl, optionally substituted

- 7 -

cycloalkyl or optionally substituted aralkyl;

$R^3 - R^6$, which can be the same or different, are hydrogen, optionally substituted (C_1-C_4) alkyl, optionally substituted aralkyl, optionally substituted phenyl or optionally substituted heteroaryl;

5 R^1 and R^2 can represent an optionally substituted heterocycle,

R^2 and R^3 can represent an optionally substituted heterocycle,

R^3 and R^5 can represent a saturated, optionally substituted carbocycle;

R^7 is hydrogen, (C_1-C_4) alkyl, (C_2-C_5) alkanoyl or aralkyl;


R^8 is (C_1-C_4) alkyl or aralkyl;

10 n is 1, 2 or 3;

X is chloride, bromide, iodide or R^2 -sulfate, wherein R^2 has the meaning indicated herein.

In one embodiment, said contacting occurs *in vitro*. In another embodiment, said contacting occurs *in vivo*.

15 The present invention is also directed to methods of treating or preventing inflammatory diseases or conditions using a hydrazino compound of Formula I or II:



20 as a racemate or an isomer, or a pharmaceutically acceptable solvate, hydrate, or salt thereof, wherein:

R^1 is hydrogen, or (C_1-C_4) alkyl, aralkyl, (C_2-C_5) alkanoyl, aroyl or heteroaroyl

R^2 is hydrogen, or optionally substituted (C_1-C_4) alkyl, optionally substituted cycloalkyl or optionally substituted aralkyl;

$R^3 - R^6$, which can be the same or different, are hydrogen, optionally

25 substituted (C_1-C_4) alkyl, optionally substituted aralkyl, optionally substituted phenyl or optionally substituted heteroaryl;

- 8 -

- R¹ and R² can represent an optionally substituted heterocycle,
R² and R³ can represent an optionally substituted heterocycle,
R³ and R⁵ can represent a saturated, optionally substituted carbocycle;
R⁷ is hydrogen, (C₁-C₄)alkyl, (C₂-C₅)alkanoyl or aralkyl;
5 R⁸ is (C₁-C₄)alkyl or aralkyl;
n is 1, 2 or 3;
X is chloride, bromide, iodide or R²-sulfate, wherein R² has the meaning indicated herein.

In one embodiment, the hydrazino compounds of Formula I or II are
10 used to treat or prevent connective tissue inflammatory conditions and diseases. In particular, the compounds can be used to treat such conditions or diseases as rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, and osteoarthritis.

In another embodiment, the compounds of Formula I or II are used to treat
15 or prevent gastrointestinal inflammatory conditions and diseases, in particular those such as Crohn's disease, ulcerative colitis, and irritable bowel syndrome.

In yet another embodiment, the hydrazino compounds of Formula I or II are used to treat central nervous system inflammatory conditions and diseases, including multiple sclerosis, Alzheimer's disease, and ischaemia-reperfusion injury
20 associated with ischemic stroke.

In another embodiment, the hydrazino compounds of Formula I or II are used to treat or prevent pulmonary inflammatory conditions and diseases. In particular, the compounds can be used to treat or prevent such conditions or diseases as asthma and adult respiratory distress syndrome.

25 In another embodiment, the hydrazino compounds of Formula I or II are used to treat or prevent chronic inflammatory skin conditions, especially such inflammatory skin conditions as psoriasis, allergic lesions, lichen planus, and pityriasis rosea.

In yet another embodiment, the hydrazino compounds of Formula I or II
30 are used to treat or prevent diseases related to carbohydrate metabolism and complications thereof, such as diabetes and complications from diabetes,

- 9 -

microvascular and macrovascular diseases such as atherosclerosis, vascular retinopathies, nephropathies and neuropathies such as polyneuropathy, mononeuropathies, and autonomic neuropathy.

In still another embodiment, the hydrazino compounds of Formula I or II
5 are used to treat or prevent diseases related to or caused by aberrations in adipocyte differentiation or function, such as atherosclerosis or obesity.

In another embodiment, the hydrazino compounds of Formula I or II are used to treat or prevent diseases related to or caused by aberrations in smooth muscle cell function, such as atherosclerosis.

10 In another embodiment, the hydrazino compounds of Formula I or II are used to treat or prevent vascular diseases, such as atheromatous and nonatheromatous arteriosclerosis, ischemic heart disease, and Raynaud's Disease and Phenomenon.

Preferred compounds are those of Formula I or II wherein n is 1. Also
15 preferred are the compounds wherein R^1 is hydrogen, and/or R^2 is hydrogen, or C_1 - C_4 alkyl, or benzyl, any of which may be optionally substituted. Preferred substituents for the benzyl group of R^2 are lower alkyl, especially methyl, and nitro, lower alkoxy, especially methoxy and halogen, especially chlorine. Especially preferred embodiments of substituted benzyl groups are *p*-toluyl, *p*-
20 nitrobenzyl, *p*-methoxybenzyl, and *p*-chlorobenzyl. Suitable values of R^2 include hydrogen, methyl, ethyl, propyl, isopropyl, benzyl, *p*-nitrobenzyl, *p*-methoxybenzyl, and *p*-chlorobenzyl. According to an embodiment, alkyl can be substituted with cycloalkyl containing 3 to 9, preferably 3 to 6 carbon atoms.

Preferred compounds of Formula I or II also include those compounds
25 wherein R^3 , R^4 , R^5 and R^6 , which can be the same or different, are hydrogen, optionally substituted C_1 - C_4 alkyl, or optionally substituted phenyl, especially when n is 1. Preferred values for R^3 , R^4 , R^5 and R^6 are hydrogen and optionally substituted phenyl. Preferred substituted phenyl groups are those substituted with a lower alkyl, especially methyl, alkoxy, such as methoxy, or a halogen such as
30 chlorine or fluorine. Especially preferred substituted phenyl groups include *o*-tolyl, *m*-tolyl, *p*-tolyl, *p*-fluorophenyl and *p*-chlorophenyl.

- 10 -

According to a preferred embodiment, one of R^5 and R^6 is hydrogen, and the other is optionally substituted phenyl, and n is preferably 1.

Another group of preferred compounds of Formula I or II are those wherein n is 1 and R^1 and R^2 or R^2 and R^3 taken together to form an optionally substituted 5-12 membered heterocyclic ring, preferably a 5 to 7 membered single ring or such a ring to which further rings are condensed (*i.e.*, a ring system). Said 5 to 7 membered ring can be either spiro or fused. The ring can be saturated or comprise double bonds. The ring or ring system can be unsubstituted or substituted, wherein the substituent can be lower alkyl, especially methyl, and nitro, lower alkoxy, especially methoxy and halogen, especially chlorine.

According to one embodiment, n is 1 and R^3 and R^5 together form a saturated carbocyclic group, which can be substituted as defined for the heterocycles above. Suitable rings include cyclopentane, cyclohexane, 4-methylcyclohexane, cycloheptane or a ring included in the adamantane ring system.

In the case where two of the substituents R^1 , R^2 , R^3 and R^5 form a ring, as described above, then it is preferred that the two remaining substituents are hydrogen. When $n > 1$, R^2 and R^3 , or R^3 and R^5 , when forming a ring, are typically on neighbouring carbon atoms. Preferably, when $n > 1$, there is at most one group R^3 and/or R^4 in the molecule different from hydrogen, such group(s) being preferably adjacent the carbon carrying R^2 .

R^1 and R^2 together are preferably saturated heterocycles, *e.g.* pyrazolidine, hexahydropyridazine and 1,2,3,4-tetrahydrophtalazine.


According to another embodiment n is 1 and the heterocyclic ring formed by the substituents R^2 and R^3 is a 5 to 6 membered nitrogen containing saturated ring. Said ring can be unsubstituted or substituted. According to a preferred embodiment the substituent is alkyl or alkoxy. According to another embodiment, this 5 to 6 membered nitrogen containing ring can be condensed another ring to form a 1,2,3,4-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline or 2,3-dihydroindole structure. As particularly preferred embodiments can be mentioned

- 11 -

piperidine, 1,2,3,4-tetrahydroisoquinoline and 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline.

In the case where R^2 and R^3 together form a heterocyclic ring, then it is preferred that R^4 is hydrogen.

5 A preferred subgenus of compounds has Formula Ia or IIa:



as a racemate or an isomer, or a pharmaceutically acceptable solvate, hydrate, or salt thereof, wherein:

n is 1,

R^1 is hydrogen,

10 R^2 is hydrogen, C_1 - C_4 alkyl or phenyl(C_1 - C_3)alkyl;

R^3 is hydrogen, C_1 - C_4 alkyl or optionally substituted phenyl; or

R^5 is hydrogen, C_1 - C_4 alkyl or optionally substituted phenyl; or

R^3 and R^5 are taken together with the carbon atoms to which they are attached to form a five to seven membered cycloalkyl ring;

15 R^4 and R^6 are independently hydrogen or C_1 - C_4 alkyl; and

R^7 is hydrogen

R^8 is C_1 - C_4 alkyl, and

X is chloride, bromide, iodide.

Examples of compounds of, and useful in, the present invention include:

20 (1*R*,2*S*)-2-(1-Methylhydrazino)-1-phenyl-1-propanol hydrogenmaleate

(1*R**,2*S**)-2-(1-Methylhydrazino)-1-phenyl-1-propanol hydrochloride

(1*R**,2*S**)-1-(2-hydroxy-1-methyl-2-phenylethyl)-1,1-dimethylhydrazinium iodide

(1*R**,2*S**)-2-(1-methylhydrazino)-1,2-diphenylethanol hydrogenmaleate

25 2-(1-methylhydrazino)-1-phenylethanol hydrogenmaleate

2-(1-methylhydrazino)-2-phenylethanol hydrogenmaleate

- 12 -

- 1-[2-methoxy-2-(*m*-methoxyphenyl)ethyl]-1-methylhydrazine hydrogenfumarate
2-Amino-6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline-1-methanol
hydrochloride
2-(1-methylhydrazino)-1-(*p*-methoxyphenyl)ethanol hydrogenfumarate
5 or a pharmaceutically acceptable salt thereof.

C₁-C₄ alkyl, as such or as a part of alkoxy or alkanoyl, is a straight or branched alkyl and thus can include methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, *tert*-butyl and isobutyl.

- 10 The term "aralkyl" as employed herein should be interpreted as any aryl attached to the alkyl, which is a chain of 1 to 6 carbon atoms and which in turn can be straight or branched. Preferably, the chain contains 1 to 3 carbon atoms. Aryl can be an monocyclic or bicyclic aromatic group containing from 6 to 12 carbons in the ring portion, preferably 6-10 carbons in the ring portion, such as
15 phenyl, naphthyl or tetrahydronaphthyl. A preferred aryl group is phenyl, which can be substituted or unsubstituted. Preferable substituents are lower alkyl (*i.e.*, C₁-C₄ alkyl), especially methyl, or a halogen or lower alkoxy, such as methoxy, or nitro. As particular preferred embodiments can be mentioned benzyl, *p*-methylbenzyl, *p*-chlorobenzyl, 2-phenylethyl and 3-phenylpropyl.

- 20 The term "(C₂-C₃)-alkanoyl" as employed herein refers to a carbonyl moiety to which is attached an alkyl group, such as any of the above C₁-C₄ alkyl groups. For example, this term includes, but is not limited to, ethanoyl, propanoyl, butanoyl, 2-methyl propanoyl.

- The term "heterocyclic ring" as used herein represents a stable 5- to 7-
25 membered mono- or bicyclic or stable 7- to 12-membered bicyclic heterocyclic ring system any ring of which may be saturated or unsaturated, and which consists of carbon atoms and from one to three heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be
30 quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. Especially useful are rings containing

- 13 -

one oxygen or sulfur, one to three nitrogen atoms, or one oxygen or sulfur combined with one or two nitrogen atoms. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic groups include piperidinyl, piperazinyl, 5 N-benzylpiperidine, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, pyrrolyl, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, 10 indolyl, quinolinyl, isoquinolinyl, chromanyl, benzimidazolyl, thiadiazoyl, benzopyranyl, benzothiazolyl, benzo[b]thiophenyl, benzo[2,3-c]1,2,5-oxadiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzothienyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl. Morpholino is the same as morpholinyl.

15 A preferred heterocyclic ring is a saturated heterocyclic ring, for example pyrrolidine, piperidine, or 1,2,3,4-tetrahydroisoquinoline, 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline, pyrazolidine, tetrahydropyridazine.

The term "heteroaryl" as employed herein refers to groups having 5 to 14 ring atoms; 6, 10 or 14 Π electrons shared in a cyclic array; and containing carbon 20 atoms and 1, 2 or 3 oxygen, nitrogen or sulfur heteroatoms (where examples of heteroaryl groups are: thienyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, furyl, pyranyl, isobenzofuranyl, benzoxazolyl, chromenyl, xanthenyl, phenoxathiinyl, 2*H*-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, 25 purinyl, 4*H*-quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinazolinyl, cinnoliny, pteridinyl, 4 α *H*-carbazolyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl and phenoxazinyl groups).

Illustrative groups include 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furyl, 3-furyl, 30 1-thienyl, 2-thienyl.

- 14 -

The term "heteroaroyl" as employed herein refers to a carbonyl moiety attached to a heteroaryl group as defined above.

The term "aroyl" as employed herein refers to a carbonyl moiety attached to an aryl group

5 The term "halogen" or "halo" as employed herein by itself or as part of another group refers to chlorine, bromine, fluorine or iodine.

 The term "substituted", unless otherwise provided for herein, refers to one or more groups independently selected from the group consisting of halo, halo (C₁₋₆)alkyl, ar(C₁₋₆)alkyl, aryl, nitro, C₁₋₆ alkoxy, and C₁₋₆ alkyl as long as the resulting
10 compound is stable. Preferred optional substituents include: halo, ar(C₁₋₆)alkyl, aryl, and C₁₋₆ alkyl.

 The term "cycloalkyl" as employed herein by itself or as part of another group refers to cycloalkyl groups containing 3 to 9 carbon atoms. Typical examples are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl,
15 cyclooctyl and cyclononyl.

 The term "haloalkyl" as employed herein refers to any of the above alkyl groups substituted by one or more chlorine, bromine, fluorine or iodine with fluorine and chlorine being preferred, such as chloromethyl, iodomethyl, trifluoromethyl, 2,2,2-trifluoroethyl, and 2-chloroethyl.

20 Some of the compounds disclosed herein may contain one or more asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms. The present invention is also meant to encompass racemic mixtures, resolved forms and mixtures thereof, as well as the individual enantiomers that may be separated according to methods that are well known to
25 those of ordinary skill in the art. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended to include both E and Z geometric isomers.

 As used herein, the term "stereoisomers" is a general term for all isomers of individual molecules that differ only in the orientation of their atoms in space.
30 It includes enantiomers and isomers of compounds with more than one chiral center that are not mirror images of one another (diastereomers).

- 15 -

The term "asymmetric center" or "chiral center" refers to a carbon atom to which four different groups are attached.

The term "enantiomer" or "enantiomeric" refers to a molecule that is nonsuperimposeable on its mirror image and hence optically active wherein the
5 enantiomer rotates the plane of polarized light in one direction and its mirror image rotates the plane of polarized light in the opposite direction.

The term "racemic" refers to a mixture of equal parts of enantiomers and which is optically inactive.


The term "resolution" refers to the separation or concentration or
10 depletion of one of the two enantiomeric forms of a molecule. The phrase "enantiomeric excess" refers to a mixture wherein one enantiomer is present in a greater concentration than its mirror image molecule.

When any variable occurs more than one time in any constituent or in Formula I or II, its definition on each occurrence is independent of its definition at
15 every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

The compounds of the present invention can be prepared by one of the following routes.

20 Compounds I were synthesized starting from amino alcohols III either *via* *N*-nitroso derivatives IV or *via* oxadiazines V. Nitroso compounds IV were obtained from amino alcohols III in slightly acidic aqueous solution by using sodium nitrite (A. A. Potekhin, A. O. Safronov, *Zhur. Org. Khim.*, 1981, 17, 379-386; H. Takahashi, T. Senda, K. Higashiyama, *Chem. Pharm. Bull.*, 1991, 39,
25 836-842; J-K. Shen, H. Katayama, N. Takatsu, I. Shiro, *J. Chem. Soc. Perkin Trans. 1*, 1993, 2087-2097) or by using other well known methods of *N*-nitrosation (M. A. Zolfigol, M. H. Zebadjadian, G. Chehardoli, H. Keypour, S. Salehzadeh, M. Shamsipur, *J. Org. Chem.*, 2000, 66, 3619-3620). Reductions of nitroso compounds IV were done either in tetrahydrofuran by using lithium
30 aluminium hydride (H. Takahashi, T. Senda, K. Higashiyama, *Chem. Pharm.*

- Bull.*, 1991, 39, 836-842) or in aqueous acetic acid by using zinc dust (D. L. Trepanier, V. Sprancmanis, K. G. Wiggs, *J. Org. Chem.*, 1964, 29, 668-672). Acidic hydrolysis of oxadiazines V (R^9 and R^{10} are (C_1-C_4) alkyl groups or can together represent a 5-7-membered saturated carbocycle), obtained from amino
- 5 alcohols III and oxaziridines VI (E. Schmitz, S. Schramm, Cs. Szántay, Zs. Kardos, *Liebigs Ann. Chem.*, 1983, 1043-1046), yielded hydrazino alcohols I. Compounds II were obtained by quaternarization of hydrazino alcohols I in acetone or acetonitrile by using alkyl halogenides or sulfates. Compounds I ($R^2 = H$) were prepared by the hydrazinolysis of the corresponding oxirane derivatives
- 10 VII. Regioisomers that formed in the oxirane ring-opening were separated by fractional crystallization, or by chromatography (M. Kim, J. D. White, *J. Am. Chem. Soc.*, 1977, 99, 1172-1180; T. Okawara, S. Ehara, H. Kagotani, Y. Okamoto, M. Eto, K. Harano, T. Yamasaki, M. Furukawa, *J. Org. Chem.*, 1996, 61, 4125-4129; T. Taguchi, J. Ishibashi, T. Matsuo, M. Kojima, *J. Org. Chem.*,
- 15 1964, 29, 1097-1103).



In case of $R^3 \neq R^4$ and $R^5 \neq R^6$, amino alcohols III were used as single

- 17 -

diastereomers. The synthesis of the enantiomers of compounds I and II started from enantiomerically pure amino alcohols III or epoxides VII. Transformations occurred without remarkable racemization.

The compounds I of this invention are useful in the form of acid addition salts. The expression "pharmaceutically acceptable acid addition salt" is intended to apply to any non-toxic organic and inorganic acid addition salts of the base compounds of Formula I and II. Illustrative inorganic acids, which form suitable salts include hydrochloric, hydrobromic, sulfuric and phosphoric acids. Illustrative organic acids, which form suitable salts include acetic, lactic, malonic, succinic, glutaric, fumaric, malic, tartaric, citric, ascorbic, maleic, benzoic, phenylacetic, cinnamic, methanesulfonic and salicylic acids.

The present invention provides a method of treating diseases in which VAP-1 has a role by selectively inhibiting VAP-1 SSAO activity, which method comprises administering to an animal in need thereof a therapeutically effective amount of a compound selected from the class of compounds depicted by Formula I or II, wherein one or more compounds of Formula I or II is administered in association with one or more non-toxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients.

The compounds of the present invention can be used to treat inflammatory conditions and diseases including but not limited to connective tissue inflammatory conditions and diseases such as ankylosing spondylitis, Reiter's syndrome, psoriatic arthritis, osteoarthritis or degenerative joint disease, rheumatoid arthritis, Sjögren's syndrome, Behçet's syndrome, relapsing polychondritis, systemic lupus erythematosus, discoid lupus erythematosus, systemic sclerosis, eosinophilic fasciitis, polymyositis and dermatomyositis, polymyalgia rheumatica, vasculitis, temporal arteritis, polyarteritis nodosa, Wegener's granulomatosis, mixed connective tissue disease, and juvenile rheumatoid arthritis; gastrointestinal inflammatory conditions and diseases such as Crohn's disease, ulcerative colitis, irritable bowel syndrome (spastic colon), fibrotic conditions of the liver, inflammation of the oral mucosa (stomatitis), and recurrent aphthous stomatitis;

- 18 -

central nervous system inflammatory conditions and diseases such as multiple sclerosis, Alzheimer's disease, and ischaemia-reperfusion injury associated with ischemic stroke; pulmonary inflammatory conditions and diseases such as asthma, chronic obstructive pulmonary disease, and adult respiratory distress syndrome;
5 and skin inflammatory conditions and diseases such as contact dermatitis, atopic dermatitis, psoriasis, pityriasis rosea, lichen planus, and pityriasis rubra pilaris.

Moreover, the compounds of the invention can be used to treat diseases related to carbohydrate metabolism and complications thereof, such as diabetes and complications of diabetes including, but not limited to microvascular and
10 macrovascular disease such as atherosclerosis, vascular retinopathies, retinopathy, nephropathy and nephrotic syndrome, neuropathies such as polyneuropathy, mononeuropathies, and autonomic neuropathy, and foot ulcers and joint problems, as well as increased risk of infection; diseases related to or caused by aberrations in adipocyte differentiation or function such as atherosclerosis and obesity; and
15 vascular diseases such as atheromatous and nonatheromatous arteriosclerosis, ischemic heart disease including myocardial infarction, peripheral arterial occlusion, thromboangiitis obliterans (Buerger's disease), and Raynaud's disease and phenomenon.

In particular, the present compounds can be used to treat atherosclerosis.
20 It is known that VAP-1 is expressed on adipocytes, smooth muscle cells, endothelial cells and is related to inflammation. Atherosclerotic plaque consists of accumulated intracellular and extracellular lipids, smooth muscle cells, connective tissue, and glycosaminoglycans. The earliest detectable lesion of atherosclerosis is the fatty streak (consisting of lipid-laden foam cells, which are macrophages that
25 have migrated as monocytes from the circulation into the subendothelial layer of the intima), which later evolves into the fibrous plaque (consisting of intimal smooth muscle cells surrounded by connective tissue and intracellular and extracellular lipids).

The term "treat inflammation" is intended to include the administration of
30 compounds of the present invention to a subject for purposes, which can include prophylaxis, amelioration, prevention or cure of an inflammatory condition or

- 19 -

disease. Such treatment need not necessarily completely ameliorate the inflammatory condition or disease. Further, such treatment can be used in conjunction with other traditional treatments for reducing the inflammatory condition known to those of skill in the art.

5 The compounds of the present invention may be administered in an effective amount within the dosage range of about 0.1 $\mu\text{g/kg}$ to about 300 mg/kg , preferably between 1.0 $\mu\text{g/kg}$ to 10 mg/kg body weight. Compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily.

10 The pharmaceutical compositions of the present invention can be administered to any animal that can experience the beneficial effects of the compounds of the invention. Foremost among such animals are humans, although the invention is not intended to be so limited.

 The pharmaceutical compositions of the present invention can be
15 administered by any means that achieve their intended purpose. For example, administration can be by parenteral, subcutaneous, intravenous, intraarticular, intrathecal, intramuscular, intraperitoneal, or intradermal injections, or by transdermal, buccal, oromucosal, ocular routes or via inhalation. Alternatively, or concurrently, administration can be by the oral route. Particularly preferred is oral
20 administration. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

 In addition to the pharmacologically active compounds, the pharmaceutical preparations of the compounds can contain suitable pharmaceutically acceptable
25 carriers comprising excipients and auxiliaries that facilitate processing of the active compounds into preparations that can be used pharmaceutically. The pharmaceutical preparations of the present invention are manufactured in a manner that is, itself, known, for example, by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus, pharmaceutical
30 preparations for oral use can be obtained by combining the active compounds with

- 20 -

solid excipients, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

Suitable excipients are, in particular, fillers such as saccharides, for example, lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example, tricalcium phosphate or calcium hydrogen phosphate, as well as binders, such as starch paste, using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents can be added, such as the above-mentioned starches and also carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings, that, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions can be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol, and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate, are used. Slow-release and prolonged-release formulations may be used with particular excipients such as methacrylic acid - ethylacrylate copolymers, methacrylic acid - ethyl acrylate copolymers, methacrylic acid - methyl methacrylate copolymers and methacrylic acid - methyl methylacrylate copolymers. Dye stuffs or pigments can be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

Other pharmaceutical preparations that can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules can contain the

- 21 -

active compounds in the form of granules that may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids such as fatty oils or liquid paraffin. In addition, stabilizers may be added.

Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example water-soluble salts and alkaline solutions. Especially preferred salts are maleate, fumarate, succinate, S,S tartrate, or R,R tartrate. In addition, suspensions of the active compounds as appropriate oily injection suspensions can be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides or polyethylene glycol-400 (the compounds are soluble in PEG-400). Aqueous injection suspensions can contain substances that increase the viscosity of the suspension, for example sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

The following examples are illustrative, but not limiting, of the method and compositions of the present invention. Other suitable modifications and adaptations of the variety of conditions and parameters normally encountered and obvious to those skilled in the art are within the spirit and scope of the invention.

Example 1

(1*R*,2*S*)-2-(1-Methylhydrazino)-1-phenyl-1-propanol hydrogenmaleate (1)

A solution of NaNO₂ (1.38 g, 20 mmol) in H₂O (10 ml) was added dropwise to a suspension of (1*R*,2*S*)-2-methylamino-1-phenyl-1-propanol hydrochloride (2.02 g, 10 mmol) in H₂O (50 ml) with vigorous stirring on an ice-cold bath, and then AcOH (0.30 g, 5 mmol) was added dropwise. The mixture was stirred at room temperature for 8 h, then was extracted with EtOAc (4 x 50 ml). The combined

- 22 -

organic phases were dried (Na_2SO_4) and evaporated under reduced pressure to give 1.86 g *N*-nitroso derivative as oily product which was used in the next step without further purification.

A solution of (1*R*,2*S*)-2-methylamino-*N*-nitroso-1-phenyl-1-propanol (1.86 g, 9.6 mmol) in THF (20 ml) was added dropwise to a stirred suspension of LiAlH_4 (0.73 g, 19.2 mmol) in THF (50 ml), and the mixture was stirred and refluxed for 3 h. The excess of LiAlH_4 was decomposed with a mixture of H_2O (1.5 ml) and THF (20 ml), the resulting precipitate was filtered off and washed with EtOAc (2 x 75 ml). The combined filtrates were dried (sicc. Na_2SO_4) and evaporated under reduced pressure. The semisolid residue was treated with an equivalent amount of maleic acid in a mixture of EtOH and Et₂O to give crystalline hydrogenmaleate salt which was filtered off and recrystallized.

¹H-NMR (400 MHz, D₂O) δ (ppm): 1.19 (3H, d, *J* = 6.8 Hz, CHCH₃), 3.09 (3H, s, NCH₃), 3.66 (1H, m, NCH), 5.40 (1H, m, OCH), 6.30 (2H, s, CHCOOH) 7.45 (5H, m, C₆H₅).

Example 2

(1*R**,2*S**)-2-(1-Methylhydrazino)-1-phenyl-1-propanol hydrochloride (2)

To a solution of 1-oxa-2-azaspiro[2.5]octane (0.60 g, 5.3 mmol) in ether (20 ml) a solution of (1*R**,2*S**)-2-methylamino-1-phenyl-1-propanol (0.86 g, 5.3 mmol) in ether (5 ml) was added. The reaction mixture was stirred at room temperature for 30 minutes then evaporated to dryness. 5% Hydrochloric acid (30 ml) was added to the residue and the mixture was stirred at ambient temperature for 1 h. The mixture was washed with Et₂O (2 x 30 ml), made alkaline with Na_2CO_3 under ice-cooling and extracted with EtOAc (3 x 50 ml). The combined EtOAc extracts were dried and evaporated to give a solide residue which was dissolved in methanol (5 ml) and converted to the crystalline hydrochloride salt by using 22 % ethanolic hydrogen chloride (2 ml) and diethyl ether. Crystals were filtered off and recrystallized from methanol/diethyl ether.

¹H-NMR (400 MHz, D₂O) δ (ppm): 1.20 (3H, d, *J* = 6.8 Hz, CHCH₃), 3.09 (3H,

- 23 -

s, NCH_3), 3.67 (1H, m, NCH), 5.41 (1H, br s, OCH), 7.47 (5H, m, C_6H_5).

Example 3

(1*R**,2*S**)-2-(1-Methylhydrazino)-1-phenyl-1-propanol hydrochloride (2)

- 5 To an ice-cooled and stirred suspension of zinc dust (2.62 g, 40 mmol) in H_2O (10 ml) a solution of (1*R**,2*S**)-2-(1-methylhydrazino)-*N*-nitroso-1-phenyl-1-propanol (1.94 g, 10 mmol, prepared according to Example 1 starting from (1*R**,2*S**)-2-methylamino-1-phenyl-1-propanol hydrochloride) in AcOH (18 ml) was added dropwise over a period of 45 min. During the addition, the temperature of the
- 10 reaction mixture was maintained at 20-25 °C by external cooling. After the addition was completed, the mixture was stirred at 50 °C for 1 h, then filtered by suction, and the zinc residue was washed with a mixture of H_2O (15 ml) and AcOH (5 ml). The combined filtrate and washings were concentrated to *ca.* 10 ml *in vacuo*. The iced-cooled solution was made basic with NaOH -solution and
- 15 extracted with Et_2O (4 x 50 ml). The combined ethereal extracts were dried and evaporated to give a yellow oil which was dissolved in methanol (5 ml) and converted to the crystalline hydrochloride salt by using 22% ethanolic hydrogen chloride (2 ml) and diethyl ether. Crystals were filtered off and recrystallized from methanol/diethyl ether.
- 20 ^1H -NMR (400 MHz, D_2O): see Example 2

Example 4

- (1*R**,2*S**)-1-(2-hydroxy-1-methyl-2-phenylethyl)-1,1-dimethylhydrazinium
- 25 iodide (4)

To a solution of (1*R**,2*S**)-2-(1-methylhydrazino)-1-phenyl-1-propanol (1.05 g, 5.8 mmol, prepared according to Example 1 starting from (1*R**,2*S**)-2-methylamino-1-phenyl-1-propanol hydrochloride) in CH_3CN (15 ml) MeI (1 ml, 16 mmol) was added. The mixture was left to stand in a closed flask at room

- 24 -

temperature for 24 h. The separated crystals were filtered off, washed with CH₃CN and recrystallized.

¹H-NMR (400 MHz, DMSO-d₆) δ (ppm): 1.16 (3H, d, *J* = 6.8 Hz, CHCH₃), 3.31 (3H, s, NCH₃), 3.32 (3H, s, NCH₃), 3.69 (1H, m, NCH), 5.61 (1H, m, OCH),
5 7.30 (1H, m, C₆H₅) 7.41 (4H, m, C₆H₅).

Example 5

(1*R**,2*S**)-2-(1-methylhydrazino)-1,2-diphenylethanol hydrogenmaleate (5)

10 A solution of NaNO₂ (1.38 g, 20 mmol) in H₂O (10 ml) was added dropwise to a suspension of (1*R**,2*S**)-2-(1-methylamino)-1,2-diphenylethanol (2.27 g, 10 mmol) in H₂O (30 ml) with vigorous stirring on an ice-cold bath, and then AcOH (0.90 g, 15 mmol) was added dropwise. The mixture was stirred at room temperature for 8 h, then was extracted with EtOAc (4 x 50 ml). The combined
15 organic phases were dried (Na₂SO₄) and evaporated under reduced pressure to give 2.04 g crystalline *N*-nitroso derivative which was used in the next step without further purification.

(1*R**,2*S**)-2-(1-methylamino)-*N*-nitroso-1,2-diphenylethanol (2.04 g, 8.0 mmol) was added in small portions to a stirred and ice-cooled suspension of LiAlH₄ (0.61 g, 16.1 mmol) in THF (80 ml), and the mixture was stirred at ambient
20 temperature for 3 h. The excess of LiAlH₄ was decomposed with a mixture of H₂O (1.2 ml) and THF (20 ml), the resulting precipitate was filtered off and washed with EtOAc (2 x 75 ml). The combined filtrates were dried (sicc. Na₂SO₄) and evaporated under reduced pressure. The oily product was treated with an
25 equivalent amount of maleic acid in a mixture of EtOH and Et₂O to give crystalline hydrogenmaleate salt which was filtered off and recrystallized.

¹H-NMR (400 MHz, D₂O) δ (ppm): 2.95 (3H, s, NCH₃), 4.45 (1H, d, *J* = 5.2 Hz, NCH₃), 5.66 (1H, d, *J* = 5.2 Hz, NCH₃), 6.30 (2H, s, CHCOOH) 7.15-7.45 (10H, om, 2 x C₆H₅).

Example 6

2-(1-methylhydrazino)-1-phenylethanol hydrogenmaleate (6) and
2-(1-methylhydrazino)-2-phenylethanol hydrogenmaleate (6a)

- 5 To a stirred solution of hydrazine hydrate (12.5 g, 0.25 mol) in EtOH (10 ml) styrene oxide (3.00 g, 25 mmol) was added dropwise. The exothermic reaction brought the mixture to reflux. After addition was complete, the mixture was kept at 60 °C for 10 min. The solvent was evaporated off and the oily residue was dissolved in EtOH and treated with an equivalent amount of maleic acid. The
10 separated crystals of 2-(1-methylhydrazino)-1-phenylethanol hydrogenmaleate were filtered off and recrystallized from EtOH.

¹H-NMR (400 MHz, D₂O) δ (ppm): 3.38 (2H, m, NCH₂), 5.07 (1H, m, OCH), 6.29 (2H, s, CHCOOH), 7.45 (5H, m, C₆H₅).

- The filtrate was treated with Et₂O to give crystalline 2-(1-methylhydrazino)-2-phenylethanol hydrogenmaleate which was filtered off and recrystallized twice
15 from a mixture of EtOH and Et₂O.

¹H-NMR (400 MHz, D₂O) δ (ppm): 3.91 (2H, m, OCH₂), 4.32 (1H, m, NCH), 6.30 (2H, s, CHCOOH), 7.47 (5H, m, C₆H₅).

20 **Example 7**

1-[2-methoxy-2-(*m*-methoxyphenyl)ethyl]-1-methylhydrazine hydrogenfumarate (7)

- A solution of NaNO₂ (2.81 g, 40.7 mmol) in H₂O (10 ml) was added dropwise to a solution of 2-methylamino-1-(*m*-methoxyphenyl)-1-ethanol hydrochloride (4.39 g, 20.2 mmol) in H₂O (20 ml) with vigorous stirring on an ice-cold bath, and then
25 AcOH (2.10 g, 35.0 mmol) was added dropwise. The mixture was stirred at 0 °C for 3 h, kept in refrigerator for 12 h, then diluted with an equal volume of H₂O, made alkaline with solid Na₂CO₃ and extracted with Et₂O (3 x 50 ml). The combined ethereal extracts were dried (Na₂SO₄) and evaporated under reduced
30 pressure to give 3.53 g yellow viscous oil which was used in the next step without

- 26 -

further purification.

55% Sodium hydride suspension (2.00 g, 45.9 mmol) was washed with *n*-hexane and suspended in THF (50 ml). A solution of 2-methylamino-*N*-nitroso-1-(*m*-methoxyphenyl)-1-ethanol (3.00 g, 14.3 mmol) in THF (90 ml) was degassed with N₂ flushing and added dropwise to the NaH suspension with stirring and continuous N₂ flushing at 0 °C over a period of 1 h. Stirring was continued at 0 °C for 2 h, then a solution of MeI (3.40 g, 24.0 mmol) in THF (30 ml) was added dropwise to the stirred suspension at 0 °C. The mixture was allowed to warm to room temperature and the excess of NaH decomposed by addition of MeOH. The solution evaporated to dryness, the residue was dissolved in H₂O (50 ml) and extracted with Et₂O (3 x 50 ml). The combined ethereal extracts were washed with H₂O (50 ml) then dried (Na₂SO₄) and evaporated under reduced pressure to give 2.70 g thick yellow oil which was used in the next step without further purification.

A solution of *N*-methyl-2-methoxy-2-(*m*-methoxyphenyl)-*N*-nitroso-ethylamine (2.70 g, 12.0 mmol) in THF (30 ml) was added dropwise to a stirred and ice-cooled suspension of LiAlH₄ (1.80 g, 47.4 mmol) in THF (90 ml). The mixture was stirred at 0 °C for 3 h, then allowed to warm to room temperature. The excess of LiAlH₄ was decomposed with a mixture of H₂O (3.6 ml) and THF (25 ml), the resulting precipitate was filtered off and washed with EtOAc (2 x 75 ml). The combined filtrate were dried (sicc. Na₂SO₄) and evaporated under reduced pressure. The oily residue was dissolved in EtOH (20 ml) and mixed with an equivalent amount of fumaric acid dissolved in warm EtOH (25 ml). After cooling the salt separated, filtered off and washed with EtOH.

¹H-NMR (400 MHz, DMSO-d₆) δ (ppm): 2.44 (3H, s, NCH₃), 2.60 (1H, m, NCH₂), 2.81 (1H, m, NCH₂), 3.14 (3H, s, CHOCH₃), 3.75 (3H, s, C₆H₄OCH₃), 4.45 (1H, m OCH), 6.57 (2H, s, CHCOOH), 6.86 (3H, m, C₆H₄), 7.27 (1H, m, C₆H₄).

- 27 -

Example 8

2-Amino-6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline-1-methanol
hydrochloride (8)

A solution of NaNO_2 (1.38 g, 20 mmol) in H_2O (10 ml) was added dropwise to a
5 suspension of 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline-1-methanol (2.23 g,
10 mmol) in H_2O (50 ml) with vigorous stirring on an ice-cold bath, and then
 AcOH (0.90 g, 15 mmol) was added dropwise. The mixture was stirred at room
temperature for 8 h, then was extracted with EtOAc (4 x 50 ml). The combined
organic phases were dried (Na_2SO_4) and evaporated under reduced pressure to
10 give 2.37 g crystalline *N*-nitroso derivative which was used in the next step
without further purification.

To a stirred suspension of 6,7-dimethoxy-2-nitroso-1,2,3,4-
tetrahydroisoquinoline-1-methanol (2.37 g, 9.4 mmol), zinc dust (2.46 g, 37.6
mmol) and H_2O (15 ml) glacial acetic acid (3.00 g, 50 mmol) was added dropwise
15 over a period of 1 h. During the addition, the temperature of the reaction mixture
was maintained at 25-30 °C by external cooling. Subsequently the reaction
mixture was stirred at 60 °C for 1 h, allowed to cool, and the excess zinc dust
filtered by suction and washed with H_2O (15 ml). The combined filtrate and
washings were made basic with aqueous NaOH -solution and extracted with CHCl_3
20 (4 x 50 ml). The combined organic phases were dried (sicc. Na_2SO_4) and
evaporated under reduced pressure. The solid residue was dissolved in methanol
(5 ml) and converted to the crystalline hydrochloride salt by using 22% ethanolic
hydrogen chloride (2 ml) and diethyl ether. Crystals were filtered off and
recrystallized from methanol/diethyl ether.

- 28 -

¹H-NMR (D₂O) δ (ppm): 3.08 (2H, m, CH₂CH₂N), 3.49 (1H, m, CH₂N), 3.76 (1H, m, CH₂N), 3.85 (3H, s, OCH₃), 3.86 (3H, s, OCH₃), 3.94 (1H, m, CH₂O), 4.19 (1H, dd, J = 12.8, 4.0 Hz, CH₂O), 4.47 (1H, m, CHN), 6.93 (1H, s, C₆H₂), 6.93 (1H, s, C₆H₂).

5

Example 9



2-(1-methylhydrazino)-1-(*p*-methoxyphenyl)ethanol hydrogenfumarate (9)




10



2-methylamino-*N*-nitroso-1-(*p*-methoxyphenyl)ethanol (1.15 g, 5.5 mmol, prepared from 2-methylamino-1-(*p*-methoxyphenyl)ethanol according to Example 1) was added in small portions to a stirred and ice-cooled suspension of LiAlH₄ (0.42 g, 11 mmol) in THF (40 ml), and the mixture was stirred at ambient
15 temperature for 2 h. The excess of LiAlH₄ was decomposed with a mixture of H₂O (0.8 ml) and THF (10 ml), the resulting precipitate was filtered off and washed with EtOAc (3 x 50 ml). The combined filtrate and washings were dried (sicc. Na₂SO₄) and evaporated under reduced pressure. The oily product was
20 treated with an equivalent amount of fumaric acid in a mixture of EtOH and Et₂O to give crystalline fumarate salt which was filtered off and recrystallized.




¹H-NMR (400 MHz, DMSO-d₆) δ (ppm): 2.57 (3H, s, NCH₃), 2.74 (2H, m, NCH₂), 3.73 (3H, s, OCH₃), 4.83 (1H, m, OCH), 6.57 (2H, s, CHCOOH), 6.89 (2H, d, J = 8.8 Hz, C₆H₄), 7.26 (2H, d, J = 8.4 Hz, C₆H₄).




Table 1. Physical data of the synthesized racemic compounds



Number	Structure	M.P. (°C)	Yield (%)	Formula (M.W.)	Elemental analysis (Calcd Found %)	Synthetic method
2		145-146	70	C ₁₀ H ₁₇ ClN ₂ O (216.71)	C 55.42 55.61 H 7.91 8.04 N 12.93 12.86	Example 1 Example 3
4		75-77	66	C ₁₁ H ₁₉ IN ₂ O (322.18)	C 41.01 40.72 H 5.94 6.18 N 8.69 8.55	Example 4




Number	Structure	Mp. (°C)	Yield (%)	Formula (M.wt)	Elemental analysis				Synthetic method
					Calcd./Found (%)				
					C	H	N		
5		152-154	71	C ₁₉ H ₂₂ N ₂ O ₅ (358.39)	63.68 63.44	6.19 6.03	7.82 7.70	Example 5	
6		120-125	48	C ₁₂ H ₁₆ N ₂ O ₅ (268.26)	53.73 53.50	6.01 5.83	10.44 10.28	Example 6	
8		210-215	67 49	C ₁₂ H ₁₉ ClN ₂ O ₃ (274.74)	52.46 52.13	6.97 6.85	10.20 10.01	Example 1 Example 8	




Number	Structure	Mp (°C)	Yield (%)	Formula (Mn)	Elemental analysis Calcd (%) Found (%)	Synthetic method
7		139-140	52	C ₁₅ H ₂₂ N ₂ O ₆ (326.34)	C: 55.21 H: 6.79 N: 8.58 O: 55.03 H: 6.45 N: 8.50	Example 7
9		142-144	67	C ₁₄ H ₂₀ N ₂ O ₆ (312.32)	C: 53.84 H: 6.45 N: 8.97 O: 53.66 H: 6.30 N: 9.08	Example 9




10		78-82	72	$C_{10}H_{16}N_2O$ (180.24)	66.64 66.28	8.95 8.72	15.54 15.49	Example 1
11		141-144	74	$C_{24}H_{36}N_4O_6$ (476.57)	60.49 60.37	7.61 7.44	11.76 11.60	Example 1
12		112-114	72	$C_{14}H_{20}N_2O_5$ (296.32)	56.75 56.51	6.80 6.49	9.45 9.26	Example 1

13		98-100	68	$C_{15}H_{22}N_2O_5$ (310.35)	58.05 57.69	7.15 7.03	9.03 8.74	Example 1
14		119-122	72 55	$C_{17}H_{21}ClN_2O$ (244.76)	58.89 59.11	8.65 8.54	11.45 11.48	Example 1 Example 3
15		133-135	70	$C_{17}H_{24}N_2O_5$ (336.38)	60.70 60.56	7.19 7.03	8.33 8.25	Example 1




16		183-186	73	$C_{16}H_{21}ClN_2O$ (292.80)	65.63 65.31	7.23 7.17	9.57 9.45	Example 1 Example 3
17		160-163	75	$C_{20}H_{24}N_2O_5$ (372.42)	64.50 64.28	6.50 6.59	7.52 7.40	Example 1




18		172-174	70	$C_{17}H_{23}ClN_2O_2$ (322.83)	63.25 62.99	7.18 7.06	8.68 8.53	Example 1
20		184-186	68	$C_{15}H_{22}N_2O_5$ (310.35)	58.05 57.92	7.15 6.94	9.03 8.95	Example 5
21		144-146	64	$C_{15}H_{22}N_2O_5$ (310.35)	58.05 58.31	7.15 7.04	9.03 8.97	Example 1

22		170 (decomp.)	53	$C_{15}H_{21}N_2O_5$ (310.35)	58.05 57.75	7.15 6.94	9.03 8.89	Example 6
23		132-134	68	$C_{16}H_{24}N_2O_5$ (324.37)	59.24 58.97	7.46 7.38	8.64 8.49	Example 1
24		90-92	74	$C_{16}H_{24}N_2O_5$ (324.37)	59.24 59.36	7.46 7.55	8.64 8.57	Example 1

29		127-128	70	$C_{13}H_{17}ClN_2O_5$ (316.74)	49.30 49.27	5.41 5.29	8.84 8.64	Example 5
30		86-88	72	$C_9H_{13}ClN_2O$ (200.66)	53.87 53.61	6.53 6.48	13.96 13.87	Example 5
33		130-133	49	$C_{13}H_{18}N_2O_6$ (298.29)	52.35 52.60	6.08 5.97	9.39 9.21	Example 6

34		88-90	73	$C_{14}H_{20}N_2O_6$ (312.32)	53.84 53.46	6.45 6.28	8.97 8.75	Example 5
35		117-120	47	$C_{13}H_{18}N_2O_6$ (298.29)	52.35 52.18	6.08 6.13	9.39 9.24	Example 6
36		135-140	49	$C_{10}H_{18}N_2O_5S$ (278.32)	43.15 42.96	6.52 6.47	10.07 9.92	Example 6

37		104-107	54	$C_{15}H_{22}N_2O_6$ (326.34)	55.21 55.03	6.79 6.84	8.58 8.55	Example 7
38		159-161	70	$C_{20}H_{24}N_2O_5$ (372.42)	64.50 64.29	6.50 6.34	7.52 7.48	Example 5
41		217 (subl.)	72	$C_{13}H_{21}ClN_2O$ (256.77)	60.81 60.72	8.24 7.93	10.91 10.85	Example 1

42		108-110	76	$C_{13}H_{20}N_2O$ (220.31)	70.87 70.63	9.15 9.01	12.72 12.48	Example 1
43		168-169	65	$C_{17}H_{24}N_2O_5$ (336.38)	60.70 60.47	7.19 6.88	8.33 8.14	Example 1
45		191-192	69 55	$C_{13}H_{21}ClN_2O_3$ (288.77)	54.07 53.88	7.33 7.40	9.70 9.67	Example 1 Example 8







46		195-197	66	$C_{10}H_{15}ClN_2O$ (214.69)	55.94 55.63	7.04 6.79	13.05 12.98	Example 5
47		246-250	71	$C_{12}H_{19}ClN_2O_3$ (274.74)	52.46 52.15	6.97 6.84	10.20 10.15	Example 5

Table 2. Physical data of the synthesized enantiomeric compounds

Number of compound	Chemical structure	mp (°C)	[α] _D ²⁰ (c)	Yield (%)	Molecular formula (M.W.)	Elemental analysis (%)		Sample	
						Calcd	Found		
1		111-113	-36 (MeOH, c = 0.1)	71	C ₁₄ H ₂₀ N ₂ O ₅ (296.32)	56.75	6.80	9.45	Example 1
						56.49	6.62	9.37	
48		112-114	+32 (MeOH, c = 0.1)	68	C ₁₄ H ₂₀ N ₂ O ₅ (296.32)	56.75	6.80	9.45	Example 1
						56.82	6.77	9.40	

Example	Structure	Mol. Wt.	Optical Activity	Mol. Wt.	Mol. Wt.	Mol. Wt.	Mol. Wt.
49		99-104	+43 (MeOH, c = 0.5)	52	C ₁₂ H ₁₆ N ₂ O ₅ (268.26)	53.73 53.64	10.44 10.38
50		98-103	-44 (MeOH, c = 0.5)	54	C ₁₂ H ₁₆ N ₂ O ₅ (268.26)	53.73 53.51	10.44 10.28

Example 10***In Vitro Inhibition of VAP-1 SSAO Activity***

- 5 VAP-1 SSAO activity was measured using the coupled colourimetric method essentially as described for monoamine oxidase and related enzymes (Holt, A., *et al.*, *Anal. Biochem.* 244:384-392 (1997)). Recombinant human VAP-1 SSAO expressed in Chinese Hamster Ovary (CHO) cells was used as a source of VAP-1 SSAO for activity measurements. Native CHO cells have negligible SSAO
- 10 activity. These cells and their culture have previously been described (Smith, D.J., *et al.*, *J. Exp. Med.* 188:17-27 (1998)). A cell lysate was prepared by suspending approximately 3.6×10^8 cells in 25ml lysis buffer (150mM NaCl, 10 mM Tris-Base pH 7.2, 1.5 mM MgCl₂, 1% NP40) and incubating at 4°C overnight on a rotating table. The lysate was clarified by centrifugation at 18000g for 5 min at room
- 15 temperature and the supernatant used directly in the assay. The VAP-1 SSAO assay was performed in 96 well microtitre plates as follows. To each well was added a predetermined amount of inhibitor if required. The amount of inhibitor varied in each assay but was generally at a final concentration of between 1 nM and 50µM. Controls lacked inhibitor. The inhibitor was in a total volume of 20:1 in
- 20 water. The following reagents were then added. 0.2M potassium phosphate buffer pH 7.6 to a total reaction volume of 200µl, 45 µl of freshly made chromogenic solution containing 1mM 2,4-dichlorophenol, 500 µM 4-aminoantipyrine and 4 U/ml horseradish peroxidase and an amount of CHO cell lysate containing VAP-1 SSAO that caused a change of 0.6 A₄₉₀ per h. This was within the linear response
- 25 range of the assay. The plates were incubated for 30 min at 37°C and the background absorbance measured at 490 nm using a Wallac Victor II multilabel counter. To initiate the enzyme reaction 20 µl 10mM benzylamine (final concentration = 1mM) was added and the plate incubated for 1 h at 37°C. The increase in absorbance, reflecting VAP-1 SSAO activity, was measured at 490nm.
- 30 Inhibition was presented as percent inhibition compared to control after correcting for background absorbance and IC₅₀ values calculated using GraphPad Prism.

Example 11

Comparison of VAP-1 SSAO activity versus total rat MAO activity

5 Rat MAO was prepared from rat liver by rinsing the $1\bar{g}$ liver sample several times in 14 ml KCl-EDTA-solution to remove all blood. Then 1 g liver sample was homogenised in 4 ml ice-cold potassium phosphate buffer (0.1 M, pH 7.4) with an Ultra-Turrax homogeniser (setting 11 000 rpm, 4 x 10s). After centrifugation at 500 g for 10 min at 4°C the supernatant was carefully withdrawn
10 and was centrifuged at 12 300 g for 15 min at 4°C. The supernatant was discharged and sedimented mitochondria were resuspended in 4 ml fresh phosphate buffer and centrifuged as previously. The mitochondria were suspended in 4 ml phosphate buffer and homogenized with an Ultra-Turrax homogeniser (setting 11 000 rpm, 2 x 10s). Mitochondrial preparate was aliquoted and stored
15 at -70°C. Total MAO activity was measured in a similar way as for VAP-1 SSAO except that 2,4-dichlorophenol was replaced by 1mM vanillic acid. To each well was added a predetermined amount of inhibitor if required. The amount of inhibitor varied in each assay but was generally at a final concentration of between 10 nM and 800 μ M. Controls lacked inhibitor. The inhibitor was in a total volume
20 of 20:1 in water. The following reagents were then added. 0.2 M potassium phosphate buffer pH 7.6 for a total reaction volume of 300 μ l, 50 μ l of freshly made chromogenic solution (as above) and 50 μ l of MAO preparation. The plates were incubated for 30 min at 37°C and the background absorbance measured at 490 nm using a Wallac Victor II multilabel counter. To initiate the enzyme
25 reaction 20 μ l of 5 mM tyramine (final concentration 0.5 mM) was added and the plate incubated for 1 h at 37°C. The increase in absorbance, reflecting MAO activity, was measured at 490nm. Inhibition was presented as percent inhibition compared to control after correcting for background absorbance and IC_{50} values calculated using GraphPad Prism. Clorgyline and pargyline (inhibitors of MAO-A

and -B respectively) at 0.5 μ M were added to some wells as positive controls for MAO inhibition.

The ability of compounds of Examples 1 to 9 to inhibit VAP-1 SSAO activity with specificity for VAP-1 SSAO over rat MAO is shown in Table 3. The results indicate that the compounds of the invention are specific inhibitors of human VAP-1 SSAO activity. The compounds of the present invention are therefore expected to have therapeutic utility in the treatment of diseases and conditions in which the SSAO activity of the human adhesion molecule VAP-1 plays a role.

Table 3

Potency and specificity of Examples 1 to 9

	Example	VAP-1 SSAO Inhibition		MAO Inhibition
		IC ₅₀ (μ M)	IC ₅₀ (μ M)	
15	4	>50	>500	\approx 10
	2	0.35	39	111
	5	0.41	54	132
	6	0.065	8.90	137
	7	0.41	44	107
20	1	0.22	31	141
	8	>10	78	<8
	9	0.31	36	116
	10	0.30	36	120
	11	0.17	16	94
25	12	0.29	40	138
	13	1.52	25	16
	14	129	>800	>6
	15	1.38	10	7
	16	4.00	249	62
	17	34	231	6

		47		
	18	2.60	22	9
	20	0.52	55	98
	21	0.26	41	158
	22	1.20	11	9
5	23	0.26	28	108
	24	0.35	11	32
	29	0.27	23	85
	30	0.31	21	68
	33	0.052	9.10	175
10	34	0.28	44	157
	35	0.029	6.00	207
	36	0.017	5.00	294
	37	0.42	44	105
	38	>50	>500	≈10
15	41	21.9	111	5
	42	3.70	106	29
	43	0.52	10.0	19
	45	8.10	45	6
	46	>10	614	<61
20	47	>10	1500	<150
	48	0.28	31	110
	49	0.14	9.80	70
	50	0.035	9.90	283
25				

Example 12

Inhibition of collagen-induced arthritis in mouse

The model in the literature:

- 5 Mouse collagen-induced arthritis (CIA) is a frequently used model both for studying the basic mechanisms of autoimmune arthritis and in assessing the efficacy of potential antiarthritic agents (van den Berg and Joosten, 1999 in *In Vivo Models of Inflammation* (Morgan DW and Marshall LA eds) pp 51-75, Birkhauser Verlag, Basel). Compounds acting through various mechanisms have
10 been demonstrated to be effective in the model and they include cyclooxygenase inhibitors, interleukins 4 and 10, leukotriene synthesis inhibitors and anti-TNF antibodies (Joosten *et al.*, *J. Immunol.* 159:4094-4102, 1997; van den Berg and Joosten, 1999 in *In Vivo Models of Inflammation* (Morgan DW and Marshall LA eds) pp 51-75, Birkhauser Verlag, Basel)).

15

Description of the model used:

- The study was conducted with groups of 14 mice to obtain statistically valid results. For arthritis induction DBA/1 mice (male, aged 10-12 weeks, approximate weight 25 g) were immunized with bovine type II collagen (100 µg) emulsified in
20 Freund's complete adjuvant by four subcutaneous injections in the back. At day 21, animals were boosted with an i.p. injection of 100 µg collagen type II diluted in PBS. This strain is highly susceptible to CIA induced with bovine type II collagen. After the second immunization, polyarthritis starts to develop in 1 to 2 weeks, with a disease incidence of approx. 80% at day 38 (Joosten *et al.*, *J.*
25 *Immunol.* 159:4094-4102, 1997). Arthritis development was scored from day 21 onwards. Animals were treated for 2.5 weeks starting after the second booster but before the arthritis onset (day 23). Intraperitoneal medication with example compound 9 (10 mg kg⁻¹ twice daily) was initiated at day 23 and continued until day 37.

Outcome:

A reduction in the cumulative score ($p < 0.05$ by Dunn's test following Kruskal-Wallis test) was detected.


The results are shown in Fig. 1

5

Having now fully described this invention, it will be understood to those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations, and other parameters without affecting the scope of the invention or any embodiment thereof. All patents
10 and publications cited herein are fully incorporated by reference herein in their entirety.

What Is Claimed Is:

1. Use of a hydrazino compound of Formula *I* or Formula *II*:



- 5 as a racemate or an isomer, or of a pharmaceutically acceptable solvate, hydrate, or salt thereof; wherein:
- R^1 is hydrogen, or $(\text{C}_1\text{-C}_4)$ alkyl, aralkyl, $(\text{C}_2\text{-C}_5)$ alkanoyl, aroyl or heteroaroyl
- R^2 is hydrogen, or optionally substituted $(\text{C}_1\text{-C}_4)$ alkyl, optionally substituted
- 10 cycloalkyl or optionally substituted aralkyl;
- $\text{R}^3 - \text{R}^6$, which can be the same or different, are hydrogen, optionally substituted $(\text{C}_1\text{-C}_4)$ alkyl, optionally substituted aralkyl, optionally substituted phenyl or optionally substituted heteroaryl;
- R^1 and R^2 can represent an optionally substituted heterocycle,
- 15 R^2 and R^3 can represent an optionally substituted heterocycle,
- R^3 and R^5 can represent a saturated, optionally substituted carbocycle;
- R^7 is hydrogen, $(\text{C}_1\text{-C}_4)$ alkyl, $(\text{C}_2\text{-C}_5)$ alkanoyl or aralkyl;
- R^8 is $(\text{C}_1\text{-C}_4)$ alkyl or aralkyl;
- n is 1, 2 or 3;
- 20 X is chloride, bromide, iodide or R^2 -sulfate, wherein R^2 has the meaning defined herein,
- for the manufacture of a pharmaceutical preparation for inhibiting copper-containing amine oxidase.

51

2. The use of a compound as defined in claim 1 for the manufacture of a pharmaceutical preparation for the treatment of an inflammatory disease or condition, a disease related to carbohydrate metabolism, a disease related to aberrations in adipocyte differentiation or function or smooth muscle cell function,
5 or a vascular disease.

3. The use according to claim 1 or 2, wherein n is 1.

4. The use according to any one of claims 1 to 3, wherein R¹ is
10 hydrogen.

5. The use of claim 1 or 2, wherein R² is benzyl optionally substituted with alkyl, nitro, methoxy, or halogen.

6. The use of claim 5, wherein R² is benzyl substituted at the *para*
15 position with methyl, nitro, methoxy, or chlorine.

7. The use according to claim 1 or 2, wherein R⁶ is phenyl optionally substituted with alkyl, nitro, methoxy, or halogen, and R⁵ is hydrogen.
20

8. The use according to claim 7, wherein the phenyl is substituted at the *para* position with methyl, nitro, methoxy, or chlorine.

9. The use of claim 1 or 2, wherein n is 1 and R³ and R⁵ together form
25 a cyclohexane ring.

10. The use of claim 1 or 2, wherein n is 1 and R² and R³ are taken together with the atoms to which they are attached to form a pyrrolidine, piperidine, tetrahydroisoquinoline, or pyrazolidine group which is optionally
30 substituted with alkyl, nitro, methoxy, or halogen, and R⁴ is hydrogen.

11. The use of claim 10, wherein R^2 and R^3 are taken together to form an optionally substituted heterocyclic ring selected from the group consisting of 1,2,3,4 -tetrahydroisoquinoline, piperidine, and 6,7-dimethoxy-1,2,3,4-tetrahydro-
5 isoquinoline.

12. The use of claim 1 or 2, wherein n is 1 and R^3 and/or R^4 is a (C_1 - C_4)alkyl group, preferably a methyl group.

10 13. The use of claim 2, wherein said inflammatory disease or condition is a connective tissue inflammatory disease or condition.

14. The use of claim 13, wherein said connective tissue inflammatory disease or condition is selected from the group consisting of ankylosing
15 spondylitis, Reiter's syndrome, psoriatic arthritis, osteoarthritis or degenerative joint disease, rheumatoid arthritis, Sjögren's syndrome, Behçet's syndrome, relapsing polychondritis, systemic lupus erythematosus, discoid lupus erythematosus, systemic sclerosis, eosinophilic fasciitis, polymyositis and dermatomyositis, polymyalgia rheumatica, vasculitis, temporal arteritis,
20 polyarteritis nodosa, Wegener's granulomatosis, mixed connective tissue disease, and juvenile rheumatoid arthritis.

15 15. The use of claim 2, wherein said inflammatory disease or condition is a gastrointestinal inflammatory disease or condition.

16. The use of claim 15, wherein said gastrointestinal inflammatory disease or condition is selected from the group consisting of Crohn's disease, ulcerative colitis, irritable bowel syndrome (spastic colon), fibrotic conditions of

53

the liver, inflammation of the oral mucosa (stomatitis), and recurrent aphthous stomatitis.

17. The use of claim 2, wherein said inflammatory disease or condition
5 is a central nervous system inflammatory disease or condition.

18. The use of claim 17, wherein said central nervous system
inflammatory disease or condition is selected from the group consisting of multiple
sclerosis, Alzheimer's disease, and ischaemia-reperfusion injury associated with
10 ischemic stroke.

19. The use of claim 2, wherein said inflammatory disease or condition
is a pulmonary inflammatory disease or condition.

20. The use of claim 19, wherein said pulmonary inflammatory disease
or condition is selected from the group consisting of asthma, chronic obstructive
pulmonary disease, and adult respiratory distress syndrome.

21. The use of claim 2, wherein said inflammatory disease or condition
20 is a skin inflammatory disease or condition.

22. The use of claim 21, wherein said skin inflammatory disease or
condition is selected from the group consisting of contact dermatitis, atopic
dermatitis, psoriasis, pityriasis rosea, lichen planus, and pityriasis rubra pilaris.

23. The use of claim 2, wherein said disease related to carbohydrate
metabolism is selected from the group consisting of diabetes, atherosclerosis,
vascular retinopathies, retinopathy, nephropathy, nephrotic syndrome,
polyneuropathy, mononeuropathies, autonomic neuropathy, foot ulcers, joint
25 problems, and increased risk of infection.

24. The use of claim 2, wherein said disease related to aberrations in adipocyte differentiation or function or smooth muscle cell function is selected from the group consisting of atherosclerosis and obesity.

5

25. The use of claim 2, wherein said vascular disease is selected from the group consisting of atheromatous arteriosclerosis, nonatheromatous arteriosclerosis, ischemic heart disease, peripheral arterial occlusion, thromboangiitis obliterans (Buerger's disease), and Raynaud's disease and phenomenon.

10

26. The use of claim 1 or 2, wherein said compound of Formula I is selected from the group consisting of:

(1*R*,2*S*)-2-(1-Methylhydrazino)-1-phenyl-1-propanol hydrogenmaleate

(1*R**,2*S**)-2-(1-Methylhydrazino)-1-phenyl-1-propanol hydrochloride

15 (1*R**,2*S**)-1-(2-hydroxy-1-methyl-2-phenylethyl)-1,1-dimethylhydrazinium iodide

(1*R**,2*S**)-2-(1-methylhydrazino)-1,2-diphenylethanol hydrogenmaleate

2-(1-methylhydrazino)-1-phenylethanol hydrogenmaleate

2-(1-methylhydrazino)-2-phenylethanol hydrogenmaleate

20 1-[2-methoxy-2-(*m*-methoxyphenyl)ethyl]-1-methylhydrazine hydrogenfumarate

2-Amino-6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline-1-methanol hydrochloride

2-(1-methylhydrazino)-1-(*p*-methoxyphenyl)ethanol hydrogenfumarate

25 or a pharmaceutically acceptable salt thereof.

27. A method of inhibiting a copper-containing amine oxidase, comprising contacting said amine oxidase with an inhibitory effective amount of a hydrazino compound of Formula I or II as defined in any of the claims 1 to 26.

5

28. The method of claim 27, wherein said contacting occurs *in vitro*.

29. The method of claim 27, wherein said contacting occurs *in vivo*.

10

30. A method of treating an inflammatory disease or condition, a disease related to carbohydrate metabolism, a disease related to aberrations in adipocyte differentiation or function or smooth muscle cell function, or a vascular disease, comprising administering to an animal in need of such treatment or prevention an effective amount of a hydrazino compound of Formula I or II as defined in any one of the claims 1 to 26.

15

1/1

Control= Vehicle (water), 2029-10= example compound 9 at a dose of 10 mg kg⁻¹.




Fig. 1

